



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07061 - Goat Anti-14-3-3 sigma / Stratifin Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: 14-3-3sigma, stratifin, HGNC:10773, SFN, YWHAS

Official Symbol: SFN

Accession Number(s): NP_006133.1

Human GenelD(s): [2810](#)

Non-Human GenelD(s): 55948 (mouse), 313017 (rat)

Immunogen

Peptide with sequence C-DLHTLSEDSYKDST, from the internal region of the protein sequence according to NP_006133.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx. 28kDa band observed in Human Skin lysates (calculated MW of 27.8kDa according to NP_006133.1). Recommended concentration: 0.03-0.1µg/ml.

IHC: In paraffin embedded Human Skin shows strong nuclear staining in selected cells of the basal layer. Recommended concentration, 2-4µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig

EB07061 (0.03µg/ml) staining of Human Skin lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB07061 (2µg/ml) staining of paraffin embedded Human Skin. Steamed antigen retrieval with Tris/EDTA buffer pH 9, HRP-staining.