

## **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

Enquiries:

UK

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06443 - Goat Anti-58KGolgi protein(Internal)/FTCD Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: FTCD, LCHC1, formiminotransferase cyclodeaminase,

formimidoyltransferase cyclodeaminase, LC1 autoantigen

Official Symbol: FTCD

Accession Number(s): NP\_006648.1; NP\_996848.1

Human GenelD(s): 10841

Important Comments: Variants (NP\_006648.1; NP\_996848.1) encode the same protein.

### **Immunogen**

Peptide with sequence CLREQGRGKDQPGRL, from the internal region of the protein sequence according to NP\_006648.1; NP\_996848.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 55kDa band observed in Human Liver lysates (calculated MW of 58.9kDa according to NP\_006648.1 and NP\_996848.1). Recommended concentration: 0.03-0.1µg/ml. Primary incubation was 1 hour.

**IHC:** In paraffin embedded Human Liver shows distinct staining of membranes and cytoplasm in a selection of hepatocytes. Recommended concentration: 3-5µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Pig

EB06443 (0.1μg/ml) staining of Human Liver lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.

EB06443 (3.8μg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.