

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06937 - Goat Anti-ADRB2R / ADRB2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** ADRB2, ADRB2R, adrenergic, beta-2-, receptor, surface [Homo sapiens], HGNC:286, ADRBR, B2AR, BAR, BETA2AR, beta-2 adrenergic receptor, beta-2 adrenoceptor, catecholamine receptor

**Official Symbol:** ADRB2

**Accession Number(s):** NP\_000015.1

**Human GeneID(s):** [154](#)

### Immunogen

Peptide with sequence C-HQGTVPDNDISQ, from the C Terminus of the protein sequence according to NP\_000015.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 48kDa band observed in Human Liver lysates (calculated MW of 46.6kDa according to NP\_000015.1). Recommended concentration: 0.1-0.3µg/ml.

### Species Reactivity

**Tested:** Human, Cow

**Expected from sequence similarity:** Human, Dog, Cow

### Specific Reference

**This antibody has been successfully used in Western blot in Cow:**

Rekawiecki R, Nowocin A, Kotwica J.

Relationship between concentrations of progesterone, oxytocin, noradrenaline, gene expression and protein level for their receptors in corpus luteum during estrous cycle in the cow.

Prostaglandins Other Lipid Mediat. 2010 Jun;92(1-4):13-8.

PMID: 20149890



EB06937 (0.2µg/ml) staining of Human Liver lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.