

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06637 - Goat Anti-AKR1C3 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** AKR1C3, DD3, HAKRB, HAKRe, HA1753, HSD17B5, hluPGFS, KIAA0119, aldo-keto reductase family 1, member C3 (3-alpha hydroxysteroid dehydrogenase, type II), prostaglandin F synthase, dihydrodiol dehydrogenase 3, chlordecone reductase homolog, hydroxysteroid (17-beta) dehydrogenase 5, type IIb 3-alpha hydroxysteroid dehydrogenase, trans-1,2-dihydrobenzene-1,2-diol dehydrogenase, DDX, aldo-keto reductase family 1, member C3, chlordecone reductase, dihydrodiol dehydrogenase X, type II 3a-hydroxysteroid dehydrogenase, type IIb 3-alpha hydroxysteroid dehydrogenase

**Official Symbol:** AKR1C3

**Accession Number(s):** NP\_003730.4

**Human GeneID(s):** [8644](#)

### Immunogen

Peptide with sequence CFASHPNYPYSDEY, from the C Terminus of the protein sequence according to NP\_003730.4.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 35kDa band observed in human liver and human breast lysates (calculated MW of 36.9kDa according to NP\_003730). Recommended concentration: 0.01-0.1µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human

EB06637 (0.03µg/ml) staining of human breast lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.