

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06637 - Goat Anti-AKR1C3 Antibody

Size: 100µg specific antibody in 200µl



#### **Target Protein**

Principal Names: AKR1C3, DD3, HAKRB, HAKRe, HA1753, HSD17B5, hluPGFS, KIAA0119, aldo-keto reductase family 1, member C3 (3-alpha hydroxysteroid dehydrogenase, type II), prostaglandin F synthase, dihydrodiol dehydrogenase 3, chlordecone reductase homolog, hydroxysteroid (17-beta) dehydrogenase 5, type IIb 3-alpha hydroxysteroid dehydrogenase, trans-1,2-dihydrobenzene-1,2-diol dehydrogenase, DDX, aldo-keto reductase family 1, member C3, chlordecone reductase, dihydrodiol dehydrogenase X, type II 3a-hydroxysteroid dehydrogenase, type IIb 3-alpha hydroxysteroid dehydrogenase

Official Symbol: AKR1C3

Accession Number(s): NP\_003730.4

Human GeneID(s): 8644

#### **Immunogen**

Peptide with sequence CFASHPNYPYSDEY, from the C Terminus of the protein sequence according to NP\_003730.4.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 35kDa band observed in human liver and human breast lysates (calculated MW of 36.9kDa according to NP\_003730). Recommended concentration: 0.01-0.1µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

EB06637 ( $0.03\mu g/ml$ ) staining of human breast lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.