

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06104 - Goat Anti-ANILLIN / Scraps (C Terminus) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ANLN, Scraps, ANILLIN, anillin, actin binding protein (scraps homolog, Drosophila), anillin (Drosophila Scraps homolog), actin binding protein, anillin, actin binding protein, DKFZp779A055, scra

Official Symbol: ANLN

Accession Number(s): NP_061155.2; NP_001271230.1; NP_001271231.1

Human GeneID(s): [54443](#)

Non-Human GeneID(s): 68743 (mouse)

Immunogen

Peptide with sequence WQPDACYKPIGKP, from the C Terminus of the protein sequence according to NP_061155.2; NP_001271230.1; NP_001271231.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

IHC: In paraffin embedded Human Kidney shows staining of nuclei in some cells of renal tubules.. Recommended concentration: 3-10µg/ml.

Immunofluorescence: Strong expression of the protein seen in the nucleus of U2OS cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of MCF7 cells. Recommended concentration: 10ug/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Cow

EB06104 (10µg/ml) staining of paraffin embedded Human Kidney. Microwaved antigen retrieval with Tris/EDTA buffer pH9, HRP-staining.

EB06104 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear staining. Actin filaments were stained with phalloidin (red). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

EB06104 Flow cytometric analysis of paraformaldehyde fixed MCF7 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.