

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06795 - Goat Anti-Arginase I Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ARG1, arginase, liver, HGNC:663, A-I, arginase, type I, arginase 1,

liver-type arginase, type I arginase

Official Symbol: ARG1

Accession Number(s): NP_000036.2

Human GeneID(s): 383

Non-Human GenelD(s): 11846 (mouse), 29221 (rat)

Immunogen

Peptide with sequence CFGLAREGNHKPID, from the C Terminus of the protein sequence according to NP_000036.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx. 37kDa band observed in Human, Mouse, Rat and Pig Liver lysates (calculated MW of 34.7kDa according to NP_000036.2). The observed molecular weight corresponds to earlier findings with different antibodies from other commercial sources.

Recommended concentration: 0.01- $0.03\mu g/ml$. Primary incubation was 1 hour. **IHC:** In paraffin embedded Human Kidney shows preferential staining of PCT.

Recommended concentration: 2-4µg/ml. Paraffin embedded Human Liver. Recommended concentration: 3.75µg/ml.

Species Reactivity

Tested: Human, Mouse, Rat, Pig

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

EB06795 (0.01µg/ml) staining of Human Liver lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB06795 (0.03µg/ml) staining of Pig (A), Mouse (B) and Rat (C) Liver lysates (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB06795 (4µg/ml) staining of paraffin embedded Human Kidney. Microwaved antigen retrieval with Tris/EDTA buffer pH9, HRP-staining.

EB06795 (3.75 μ g/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.