

## **UK Office**

**Everest Biotech Ltd** 

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06344 - Goat Anti-BAF53A and BAF53B Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** BAF53A, BAF53B, ACTL6, BAF53a, BAF53b, actin-like 6A, actin-like 6B, hArpN alpha, actin-related protein, ACTL6A, ACTL6B, 53 kDa BRG1-associated factor B, ARPN-BETA, Arp4, INO80K, MGC5382, BAF complex 53 kDa subunit, BAF53, BRG1-associated factor, INO80 complex subunit K, actin-related protein 4, hArpN beta **Official Symbol:** ACTL6A / ACTL6B

Accession Number(s): NP\_004292.1; NP\_817126.1; NP\_057272.1

# Human GenelD(s): <u>51412</u>, <u>86</u>

**Important Comments:** This antibody is expected to recognise both BAF53A and BAF53B, which are almost identical. Reported variants represent identical protein (NP\_817126.1; NP\_829888.1).

## Immunogen

Peptide with sequence YEEGGKQCVERKCP, from the C Terminus of the protein sequence according to NP\_004292.1; NP\_817126.1; NP\_057272.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

**Peptide ELISA:** antibody detection limit dilution 1:64000. **Western blot:** Approx 48kDa band observed in Hela lysates (predicted MW of 49kDa according to NP\_057272, NP\_004292). Recommended for use at 1-3µg/ml.

#### **Species Reactivity**

Tested: Human Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow, EB06344 staining (1µg/ml) of Hela lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.