

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05111-T - Goat Anti-BLNK / SLP-65 Antibody - Trial

Size: 20µg specific antibody in 40µl



### Target Protein

**Principal Names:** BLNK, SLP-65, B-cell linker, BASH, BLNK-s, Ly57, MGC111051, SLP65, B cell linker protein, B-cell adapter containing a SH2 domain protein, B-cell adapter containing a Src homology 2 domain protein, OTTHUMP00000020167, Src homology 2 domain-containing leukocyte protein of 65 kDa, BLNK-S, LY57, OTTHUMP00000020168

**Official Symbol:** BLNK

**Accession Number(s):** NP\_037446.1; NP\_001107566.1

**Human GeneID(s):** [29760](#)

**Non-Human GeneID(s):** 17060 (mouse)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_037446.1; NP\_001107566.1).

### Immunogen

Peptide with sequence C-KDSTRLKYAVKVS, from the C Terminus of the protein sequence according to NP\_037446.1; NP\_001107566.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 70kDa band observed in lysates of cell line Daudi (calculated MW of 50.5kDa according to NP\_037446.1). The observed molecular weight corresponds to how SLP-65 is expected to run. However, when transiently expressed in HEK293, a 50kDa band is observed. Recommended concentration: 1-3µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig, Cow

EB05111 staining (4µg/ml) of Daudi lysate (RIPA buffer, 30µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

HEK293 overexpressing BLNK (RC202488) and probed with EB05111 (mock transfection in first lane), tested by Origene.