

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08821 - Goat Anti-CACNA1C Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CACNA1C, calcium channel, voltage-dependent, L type, alpha 1C subunit, CACH2, CACN2, CACNL1A1, CCHL1A1, CaV1.2, MGC120730, TS, DHPR, alpha-1 subunit, calcium channel, L type, alpha 1 polypeptide, isoform 1, cardic muscle, calcium channel, cardic dihydropyridine-sensitive, alpha-1 subunit, voltage-dependent L-type calcium channel alpha 1C subunit, voltage-gated L-type calcium channel Cav1.2 alpha 1 subunit, splice variant 10*, voltage-gated calcium channel alpha subunit Cav1.2

Official Symbol: CACNA1C Accession Number(s): NP_000710.5; NP_955630.2; NP_001161097.1; NP_001123299.1; NP_001123301.1; NP_001161096.1; NP_001123302.1;

NP_001123303.1 and more **Human GenelD(s):** 775

Important Comments: This antibody is expected to recognise all reported human isoforms.

Immunogen

Peptide with sequence C-RARGRPSEELQD, from the C Terminus of the protein sequence according to NP_000710.5; NP_955630.2; NP_001161097.1; NP_001123299.1; NP_001123301.1; NP_001161096.1; NP_001123302.1; NP_001123303.1 and more.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx. 200kDa band observed in Human Heart lysate (calculated MW of 240kDa according to NP_000710.5). Recommended concentration: 1.0-3.0µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

EB08821 ($1\mu g/ml$) staining of Human Heart lysate ($35\mu g$ protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.