

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB05563 - Goat Anti-CAPE Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: SMC2, structural maintenance of chromosomes 2, PRO0324, CAP-E, CAPE, FLJ10093, SMC2L1, hCAP-E, OTTHUMP00000021818, OTTHUMP00000021819, SMC protein 2, SMC-2, SMC2 (structural maintenance of chromosomes 2, yeast)-like 1, SMC2 structural maintenance of chromosomes 2-like 1, XCAP-E homolog, chromosome-associated protein E, structural maintenance of chromosomes (SMC) family member, chromosome-associated protein E, structural maintenance of chromosomes

Official Symbol: SMC2

Accession Number(s): NP_006435.2

Human GenelD(s): 10592

Important Comments: Reported variants NP_006435.2, NP_001036015.1 and

NP_001036016.1 represent identical protein

Immunogen

protein 2

Peptide with sequence C-SKAKPPKGAHVEV, from the C Terminus of the protein sequence according to NP_006435.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 140kDa band observed in lysates of cell line HeLa (calculated MW of 136kDa according to NP_006435.2). Recommended concentration: 0.2-0.6µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa

20kDa

15kDa

EB05563 (0.2 μ g/ml) staining of HeLa lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.