

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07271 - Goat Anti-Chromogranin A precursor Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: chromogranin A precursor, chromogranin A (parathyroid secretory protein 1), CHGA, CGA, parathyroid secretory protein 1, Vasostatin, Pancreastatin,

Parastatin, Pituitary secretory protein I, SP1

Official Symbol: CHGA

Accession Number(s): NP_001266.1

Human GeneID(s): 1113

Immunogen

Peptide with sequence C-QGAKERAHQQKKH, from the internal region of the protein sequence according to NP_001266.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.

Western blot: Approx 50kDa band observed in Human Kidney lysates (calculated MW of 50.7 kDa according to NP_001266.1). Recommended concentration: 0.1-0.3μg/ml. IHC: Paraffin embedded Human Pancreas and Small Intestine. Recommended

concentration: 3.75µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

EB07271 (0.1μg/ml) staining of Human Kidney lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB07271 (3.75µg/ml) staining of paraffin embedded Human Pancreas. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB07271 (3.75µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.