

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB10235 - Goat Anti-CIRBP (aa 81-91) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CIRP, cold inducible RNA binding protein, cold inducible RNA-binding protein, Cold-inducible RNA-binding protein, glycine-rich RNA binding protein, CIRBP

**Official Symbol:** CIRBP

**Accession Number(s):** NP\_001271.1

**Human GeneID(s):** [1153](#)

**Non-Human GeneID(s):** 12696 (mouse), 81825 (rat)

### Immunogen

Peptide with sequence C-QAGKSSDNRSR, from the internal region of the protein sequence according to NP\_001271.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 18kDa band observed in Mouse and Rat Testis lysates (calculated MW of 18.6kDa according to Mouse NP\_031731.1 and Rat NP\_112409.2).

Recommended concentration: 0.05-0.2µg/ml. An additional band of unknown identity was also consistently observed at 23kDa. This band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

### Species Reactivity

**Tested:** Mouse, Rat

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow



EB10235 (0.05µg/ml) staining of Mouse Testis lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.