

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06929 - Goat Anti-Clusterin / APOJ Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: APOJ, clusterin, CLU, complement lysis inhibitor, SP-40,40, sulfated glycoprotein 2, testosterone-repressed prostate message 2, apolipoprotein J, HGNC:2095, CLI, MGC24903, SGP-2, SGP2, SP-40, TRPM-2, TRPM2, apolipoprotein J, clusterin, complement-associated protein SP-40, sulfated glycoprotein 2, AAG4, KUB1, aging-associated protein 4, clusterin (complement lysis inhibitor, SP-40,40, sulfated glycoprotein 2, testosterone-repressed prostate message 2, apolipoprotein J)

Official Symbol: CLU

Accession Number(s): NP_001822.2; NP_976084.1

Human GeneID(s): [1191](#)

Important Comments: This antibody is expected to recognise both reported isoforms (NP_001822.2 and NP_976084.1)

Immunogen

Peptide with sequence C-EKALQEYRKKHREE, from the C Terminus of the protein sequence according to NP_001822.2; NP_976084.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 35-37kDa band observed in Human Colon and Prostate lysates (calculated MW of 57.8kDa according to NP_NP_001822.2 and 52.5 according to NP_976084.1). The observed molecular weight corresponds to earlier findings in literature (PMID: 9713325 and PMID: 11560857). Recommended concentration: 0.03-0.1µg/ml. Primary incubation was 1 hour.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog

EB06929 (0.1µg/ml) staining of Human Colon lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB06929 (5µg/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining. **This data is from a previous batch, not on sale.**