

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08954 - Goat Anti-COG7 Antibody

Size: 100µg specific antibody in 200µl

Target Protein

Principal Names: COG7, component of oligomeric golgi complex 7, CDG2E Official Symbol: COG7 Accession Number(s): NP_705831.1 Human GeneID(s): <u>91949</u>

Immunogen

Peptide with sequence C-KTRPEDYRQVSK, from the internal region of the protein sequence according to NP_705831.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.


Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000. **Western blot:** Approx. 85kDa band observed in lysates cell lines HeLa, Jurkat and MOLT4 (calculated MW of 86.3kDa according to NP_705831.1). Recommended concentration: 0.03-0.1µg/ml. Primary incubation was 1 hour. **IHC:** Paraffin embedded Human Testis. Recommended concentration: 5µg/ml.

Species Reactivity


Tested: Human Expected from sequence similarity: Human



1	250kDa 150kDa 100kDa 75kDa
	50kDa
	37kDa
	25kDa
	20kDa

15kDa

EB08954 (0.03µg/ml) staining of MOLT4 lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB08954 (5µg/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.