

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05285-T - Goat Anti-COX1 / PTGS1 Antibody - Trial

Size: 20µg specific antibody in 40µl



Target Protein

Principal Names: PTGS1, COX1, COX3, PHS1, PCOX1, PGHS1, PTGHS, PGG/HS, PGHS-1, prostaglandin-endoperoxide synthase 1 (prostaglandin G/H synthase and cyclooxygenase), prostaglandin G/H synthase and cyclooxygenase, RP11-542K23.6

Official Symbol: PTGS1

Accession Number(s): NP_000953.2; NP_542158.1

Human GeneID(s): [5742](#)

Important Comments: This antibody is expected to recognise both human isoforms of this protein according to NP_000953.2 and NP_542158.1.

Immunogen

Peptide with sequence C-QDDGPAVERPSTEL, from the C Terminus of the protein sequence according to NP_000953.2; NP_542158.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Approx 60kDa and 70-75kDa bands observed in lysates of cell line U937 and approx 70-75kDa band observed in Mouse Kidney and Skeletal Muscle lysates (calculated MW of 64.5kDa according to Human NP_542158.1 and 68.7kDa Human NP_000953.2, and 69.0kDa according to Mouse NP_032995.1). The observed molecular weights correspond to earlier findings with different antibodies from other commercial sources. Recommended concentration: 0.5-2µg/ml.

Immunofluorescence: Strong expression of the protein seen in the Golgi apparatus and vesicles of NIH3T3 cells. Recommended concentration: 5µg/ml.

Species Reactivity

Tested: Human, Mouse


Expected from sequence similarity: Human




EB05285 (2 μ g/ml) staining of U937 lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.



EB05285 (0.5 μ g/ml) staining of Mouse Kidney (A) and Skeletal Muscle (B) lysate (35 μ g protein in RIPA buffer).
Primary incubation was 1 hour. Detected by chemiluminescence.



EB05285 Immunofluorescence analysis of paraformaldehyde fixed NIH3T3 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (5ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing Golgi and vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (5ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05285 (3.8µg/ml) staining of paraffin embedded Human Heart. Steamed antigen retrieval with citrate buffer pH 6, AP-staining. **This data is from a previous batch, not on sale.**