

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05118 - Goat Anti-CSK Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CSK, c-src tyrosine kinase, MGC117393

Official Symbol: CSK

Accession Number(s): NP_004374.1

Human GeneID(s): [1445](#)

Non-Human GeneID(s): 12988 (mouse), 315707 (rat)

Important Comments: Reported variants represent identical protein: NP_004374.1; NP_001120662.1.

Immunogen

Peptide with sequence C-EQLEHIKTHELHL, from the C Terminus of the protein sequence according to NP_004374.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 50kDa band observed in lysates of cell line Jurkat and in Mouse Spleen lysates (calculated MW of 50.7kDa according to NP_004374.1). Recommended concentration: 1-3µg/ml.

IHC: In paraffin embedded Human Colon shows staining of a cell layer between the mucosal columns and the muscularis mucosae. Recommended concentration: 3-6µg/ml.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

EB5118 (0.5µg/ml) staining of Jurkat lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.

EB05118 (1µg/ml) staining of Mouse Spleen lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.

EB05118 (4µg/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining.