

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06750 - Goat Anti-Dipeptidylpeptidase 10 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** DPP10, dipeptidylpeptidase 10, HGNC:20823, DPRP3, dipeptidyl peptidase IV-related protein 3, dipeptidyl peptidase X, dipeptidyl peptidase-like protein 2, dipeptidyl-peptidase 10, DPL2, DPPY, dipeptidyl peptidase 10, dipeptidyl peptidase like protein 2

Official Symbol: DPP10

Accession Number(s): NP\_065919.2; NP\_001004360.2

Human GeneID(s): 57628

Important Comments: This antibody is expected to recognise both reported isoforms

NP\_065919.2 and NP\_001004360.2.

### **Immunogen**

Peptide with sequence C-DEGHNVSEKSKYHL, from the internal region of the protein sequence according to NP\_065919.2; NP\_001004360.2.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 90kDa band observed in Human Bone Marrow lysates (calculated MW of 90.0kDa according to NP\_065919.2 and of 90.1 according to NP\_001004360.1). Recommended concentration: 0.1-0.3µg/ml.

Recommended concentration. 0.1-0.5µg/mi.

IHC: Paraffin embedded Human Brain (Cortex and Cerebellum). Recommended concentration: 3.75µg/ml.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Dog, Cow

EB06750 (0.1μg/ml) staining of Human Bone Marrow lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB06750 (3.75µg/ml) staining of paraffin embedded Human Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB06750 (3.75µg/ml) staining of paraffin embedded Human Cerebellum. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.