

### **UK Office**

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

#### www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06740 - Goat Anti-DOPA decarboxylase Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: dopa decarboxylase (aromatic L-amino acid decarboxylase),
HGNC:2719, DDC, AADC, aromatic L-amino acid decarboxylase
Official Symbol: DDC
Accession Number(s): NP\_000781.1; NP\_001229815.1; NP\_001229816.1;
NP\_001229817.1; NP\_001229818.1
Human GenelD(s): 1644
Important Comments: This antibody is expected to recognise isoforms 1, 2, 3, 4 and 5

(NP\_000781.1; NP\_001229815.1; NP\_001229816.1; NP\_001229817.1; NP\_001229818.1 respectively). Reported variants represent identical protein (NP\_000781.1; NP\_001076440.1).

### Immunogen

Peptide with sequence C-WEHIKELAADVL, from the C Terminus of the protein sequence according to NP\_000781.1; NP\_001229815.1; NP\_001229816.1; NP\_001229817.1; NP\_001229818.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

**Peptide ELISA:** antibody detection limit dilution 1:8000. **Western blot:** Approx 50kDa band observed in Human Kidney lysates and 48kDa on Human Brain lysates (calculated MW of 53.9kDa according to NP\_000781.1). Recommended concentration: 0.03-0.1µg/ml.

# **Species Reactivity**

Tested: Human Expected from sequence similarity: Human

	250kDa 150kDa 100kDa
-	75kDa 50kDa
ĺ.	37kDa
	25kDa
	20kDa
	15kDa
	10kDa

EB06740 (0.03µg/ml) staining of Human Kidney lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.