

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07081 - Goat Anti-ERG Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** ERG, v-ets erythroblastosis virus E26 oncogene like (avian), HGNC:3446, erg-3, p55, transcriptional regulator ERG (transforming protein ERG), v-ets avian erythroblastosis virus E26 oncogene related, v-ets erythroblastosis virus E26 oncogene like, v-ets erythroblastosis virus E26 oncogene homolog (avian), OTTHUMP00000115970, OTTHUMP00000208036, TMPRSS2-ERG prostate cancer specific, ets-related

**Official Symbol:** ERG

**Accession Number(s):** NP\_891548.1; NP\_004440.1; NP\_001129626.1; NP\_001129627.1; NP\_001230358.1

**Human GeneID(s):** [2078](#)

**Non-Human GeneID(s):** 13876 (mouse), 170909 (rat)

**Important Comments:** This antibody is expected to recognise all reported isoforms (NP\_891548.1; NP\_004440.1; NP\_001129626.1; NP\_001129627.1; NP\_001230358.1)

### Immunogen

Peptide with sequence C-PNTRLPTSHMP SH, from the C Terminus of the protein sequence according to NP\_891548.1; NP\_004440.1; NP\_001129626.1; NP\_001129627.1; NP\_001230358.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 55kDa band observed in lysates of cell line MOLT4 (calculated MW of 53.8kDa according to NP\_891548.1). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat

EB07081 (0.1µg/ml) staining of MOLT4 lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.  
EB07081 (10µg/ml) staining of paraffin embedded Human Prostate. Steamed antigen retrieval with citrate buffer pH 6, AP-staining. **This data is from a previous batch, not on sale.**