

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09167 - Goat Anti-EWS / EWSR1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: EWSR1, EWS, Ewing sarcoma breakpoint region 1, EWS/FLI fusion protein, EWSR1/ZNF384 fusion protein, Ewings sarcoma EWS-Flt1 (type 1) oncogene, bK984G1.4 (Ewing sarcoma breakpoint region 1 protein), extraskeletal myxoid chondrosarcoma EWS/TEC/CHN fusion protein

Official Symbol: EWSR1

Accession Number(s): NP_053733.2; NP_005234.1; NP_001156757.1; NP_001156758.1; NP_001156759.1

Human GeneID(s): [2130](#)

Non-Human GeneID(s): 14030 (mouse), 289752 (rat)

Important Comments: This antibody is expected to recognize all five reported isoforms (NP_053733.2; NP_005234.1; NP_001156757.1; NP_001156758.1; NP_001156759.1).

Immunogen

Peptide with sequence C-TSYDQSSYSQQNTYG, from the internal region of the protein sequence according to NP_053733.2; NP_005234.1; NP_001156757.1; NP_001156758.1; NP_001156759.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx. 90kDa band observed in nuclear lysates of cell line HeLa (calculated MW of 68.5kDa according to NP_053733.1). The observed molecular weight corresponds to earlier findings with different antibodies from other commercial sources. Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Pig, Cow

EB09167 (0.3µg/ml) staining of nuclear HeLa lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB09167 (3.75µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining. **This data is from a previous batch, not on sale.**

EB09167 (3.75µg/ml) staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining. **This data is from a previous batch, not on sale.**