

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07742-B - Goat Anti-FABP2, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** FABP2, fatty acid binding protein 2, intestinal, FABPI, I-FABP, fatty acid-binding protein 2, intestinal-type fatty acid-binding protein, Fatty acid-binding protein, intestinal, intestinal fatty acid binding protein 2, MGC133132

Official Symbol: FABP2

Accession Number(s): NP\_000125.2

Human GeneID(s): 2169

Non-Human GeneID(s): 14079 (mouse)

# **Immunogen**

Peptide with sequence C-EGVEAKRIFKKD., from the C Terminus of the protein sequence according to NP\_000125.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.

**Western blot:** Approx 16kDa band observed in Mouse Duodenum lysates (calculated MW of 15.1kDa according to Mouse NP\_032006.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

# **Species Reactivity**

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

Biotinylated EB07742 (0.1μg/ml) staining of Mouse Duodenum lysate (35μg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.