

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12554 - Goat Anti-FCER1A (aa177-189) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: FCER1A, Fc fragment of IgE, high affinity I, receptor for, alpha polypeptide, FCE1A, FcERI, Fc IgE receptor, alpha polypeptide, Fc epsilon RI alpha-chain, Fc-epsilon RI-alpha, high affinity immunoglobulin epsilon receptor alpha-subunit, high affinity immunoglobulin epsilon receptor subunit alpha, IgE Fc receptor subunit alpha, immunoglobulin E receptor, high-affinity, of mast cells, alpha polypeptide

Official Symbol: FCER1A

Accession Number(s): NP_001992.1

Human GeneID(s): [2205](#)

Immunogen

Peptide with sequence C-TGKVVQLDYESEP, from the internal region of the protein sequence according to NP_001992.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 50kDa band observed in Human Peripheral Blood Mononucleocytes lysates (calculated MW of 29.6kDa according to NP_001992.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Ganguly et al, J Biol Chem. 2007 Nov 9;282(45):32758-64.PMID: 17728245). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

EB12554 (0.3µg/ml) staining of Human Peripheral Blood Lymphocytes lysate (35µg protein in RIPA buffer).
Primary incubation was 1 hour. Detected by chemiluminescence.