

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06303 - Goat Anti-FHL2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** FHL2, DRAL, AAG11, SLIM3, four and a half LIM domains 2, down-regulated in rhabdomyosarcoma LIM protein, aging-associated gene 11, four and a half LIM-domain protein 2, OTTHUMP00000203164, OTTHUMP00000203168, OTTHUMP00000203169

**Official Symbol:** FHL2

**Accession Number(s):** NP\_001441.4

**Human GeneID(s):** [2274](#)

**Non-Human GeneID(s):** 14200 (mouse), 63839 (rat)

**Important Comments:** Reported variants represent identical protein (NP\_001441.4; NP\_963849.1; NP\_963851.2; NP\_001034581.1).

### Immunogen

Peptide with sequence RDDILCPDCGKDI, from the C Terminus of the protein sequence according to NP\_001441.4.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 30kDa band observed in Human and Mouse Heart lysates (calculated MW of 32.2kDa according to NP\_001441.4). Recommended concentration: 1-3µg/ml.

### Species Reactivity

**Tested:** Human, Mouse

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow



EB06303 (1µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.