

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07434 - Goat Anti-FLAP / ALOX5AP (internal) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** ALOX5AP, FLAP, arachidonate 5-lipoxygenase-activating protein, MK-886-binding protein, five-lipoxygenase activating protein

**Official Symbol:** ALOX5AP

**Accession Number(s):** NP\_001620.2; NP\_001191335.1

**Human GeneID(s):** [241](#)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_001620.2; NP\_001191335.1).

### Immunogen

Peptide with sequence C-HKVEHESRTQNGR, from the internal region of the protein sequence according to NP\_001620.2; NP\_001191335.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:128000.

**Western blot:** Approx 16kDa band observed in lysates of U937 and of Peripheral Blood Mononucleocytes (calculated MW of 18.2kDa according to NP\_001620.2). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

**IHC:** In paraffin embedded Human Spleen shows staining of select splenocytes. Paraffin embedded Human Lung and Small Intestine. Recommended concentration: 3.75µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Dog

EB07434 (0.1µg/ml) staining of Human U937 cell lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB07434 (3.75µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB07434 (3.75µg/ml) staining of paraffin embedded Human Lung. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB07434 (3.75µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.