

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07520 - Goat Anti-FOXP2 (internal) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** FOXP2, forkhead box P2, CAGH44, DKFZp686H1726, SPCH1, TNRC10, CAG repeat protein 44, forkhead/winged-helix transcription factor, speech and language disorder 1, trinucleotide repeat containing 10

**Official Symbol:** FOXP2

**Accession Number(s):** NP\_055306.1; NP\_683696.2; NP\_683697.2; NP\_683698.2; NP\_001166237.1

**Human GeneID(s):** [93986](#)

**Non-Human GeneID(s):** 114142 (mouse)

**Important Comments:** This antibody is expected to recognize isoform I (NP\_055306.1), isoform II (NP\_683696.2), isoform IV (NP\_683698.2) and isoform V (NP\_001166237.1).

### Immunogen

Peptide with sequence C-DEVEYQKRRSQKIT, from the internal region of the protein sequence according to NP\_055306.1; NP\_683696.2; NP\_683697.2; NP\_683698.2; NP\_001166237.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:64000.

**Western blot:** Approx 85kDa band observed in Human Brain (Cerebellum and Frontal Cortex) lysates and in Rat Brain lysates (calculated MW of 82.6kDa according to NP\_683696.1). Recommended concentration: 0.3-1µg/ml. An additional band of unknown identity was also consistently observed at 28-30kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

**IHC:** In paraffin embedded Human Prostate shows nuclear staining in the secretory cells. Recommended concentration, 2-4µg/ml.

### Species Reactivity

**Tested:** Human, Rat

**Expected from sequence similarity:** Human, Mouse, Rat, Cat, Dog, Pig, Cow, Zebrafish

### Specific Reference

Fujita H, Morita N, Furuichi T, Sugihara I.

Clustered fine compartmentalization of the mouse embryonic cerebellar cortex and its rearrangement into the postnatal striped configuration.

J Neurosci. 2012 Nov 7;32(45):15688-703.

PMID: 23136409

EB07520 (0.3µg/ml) staining of Human Brain (Cerebellum) lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB07520 (2.5µg/ml) staining of paraffin embedded Human Prostate. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.