

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08257 - Goat Anti-C2GnT-M (aa 273 to 284) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: GCNT3, glucosaminyl (N-acetyl) transferase 3, mucin type, C2/4GnT,

C2GnT-M, C2GnT2, GnT-M, beta1,6-N-acetylglucosaminyltransferase

Official Symbol: GCNT3

Accession Number(s): NP_004742.1

Human GeneID(s): 9245

Immunogen

Peptide with sequence C-RDTLHLTNKKKD, from the internal region of the protein sequence according to NP_004742.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx. 50kDa band observed in Human Lymph Node lysates (calculated MW of 50.9kDa according to NP_004742.1). In transfected HEK293 transiently expressing GCNT3 a band of approx. 55kDa is observed. This band is not observed in the non-transfected HEK293. Recommended concentration: 0.3-1µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB08257 (0.3 μ g/ml) staining of human lymph node lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



HEK293 overexpressing GCNT3 (RC202007) and probed with EB08257 (mock transfection in first lane), tested by Origene.