

## **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre
77 Heyford Park

Upper Heyford Oxfordshire

**OX25 5HD** 

UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10456-B - Goat Anti-GNB3 & GNB4, Biotinylated Antibody

Size: 100µg specific antibody in 200µl

# **Target Protein**

**Principal Names:** G protein, beta-3 subunit, GTP-binding regulatory protein beta-3 chain, guanine nucleotide binding protein (G protein), beta polypeptide 3, guanine nucleotide-binding protein G(I)/G(S)/G(T) beta subunit 3, guanine nucleotide-binding

protein, beta-3, GNB3

Official Symbol: GNB3

Accession Number(s): NP\_002066.1, NP\_067642.1

Human GenelD(s): <u>2784</u>, <u>59345</u> Non-Human GenelD(s): 60449 (rat)

Important Comments: This antibody is expected to recognize GNB3 & GNB4 (GeneID:

2784, 59345; ProteinID: NP\_002066.1; NP\_067642.1).

### **Immunogen**

Peptide with sequence SGHDNRVSCLGVT., from the internal region of the protein sequence according to NP\_002066.1, NP\_067642.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

**Western blot:** Approx 35kDa band observed in lysates of cell line HepG2 (calculated MW of 37.2kDa according to NP\_002066.1. Recommended concentration: 0.1-0.3µg/ml.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

Biotinylated EB10456 (0.1µg/ml) staining of HepG2 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.