

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD

Enquiries:

info@everestbiotech.com

Sales:

UK

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07290-T - Goat Anti-GPR94 / TRA1 Antibody - Trial

Size: 20µg specific antibody in 40µl



Target Protein

Principal Names: TRA1, GRP94, tumor rejection antigen (gp96) 1, HGNC:12028, ECGP, GP96, Tumor rejection antigen-1 (gp96), endothelial cell (HBMEC) glycoprotein, glucose

regulated protein, 94 kDa, tumor rejection antigen (gp96) 1

Official Symbol: HSP90B1

Accession Number(s): NP_003290.1

Human GeneID(s): 7184

Non-Human GeneID(s): 22027 (mouse), 362862 (rat)

Immunogen

Peptide with sequence C-KEGVKFDESEKTKE, from the internal region of the protein sequence according to NP_003290.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 100kDa band observed in lysates of cell lines NIH3T3 (calculated MW of 92.5kDa according to NP_003290.1). Recommended concentration: 0.01-0.03µg/ml.

IHC: In paraffin embedded Human Liver shows staining in some hepatocyte nuclei and weak cytoplasm staining of hepatocytes. Paraffin embedded Human Uterus.

Recommended concentration: 3-5µg/ml.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow, Zebrafish

EB07290 (0.03μg/ml) staining of NIH3T3 lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour.

Detected by chemiluminescence.

EB07290 (3.8μg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB07290 (5µg/ml) staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.