

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05284-T - Goat Anti-GRB2 Antibody - Trial

Size: 20µg specific antibody in 40µl



Target Protein

Principal Names: GRB2, growth factor receptor-bound protein 2, ASH, EGFRBP-GRB2, Grb3-3, MST084, MSTP084, HT027, abundant SRC homology, epidermal growth factor receptor-binding protein GRB2, growth factor receptor-bound protein 3, NCKAP2, protein Ash, SH2/SH3 adapter GRB2

Official Symbol: GRB2

Accession Number(s): NP_002077.1; NP_987102.1

Human GeneID(s): [2885](#)

Non-Human GeneID(s): 14784 (mouse), 81504 (rat)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_002077.1 and NP_987102.1).

Immunogen

Peptide with sequence C-PRNYVTPVNRNV, from the C Terminus of the protein sequence according to NP_002077.1; NP_987102.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 26kDa band observed in lysates of cell line MOLT4, of Human Thymus and of Human, Mouse, Rat, Pig Brain and Spleen lysates (calculated MW of 25.2kDa according to Human NP_002077.1 and according to Rat NP_110473.2). Recommended for use at 0.05-0.15µg/ml.

IHC: In paraffin embedded Human Lymph Node shows nuclear staining in some cells inside the germinal centre. Recommended concentration: 1-2µg/ml.

Species Reactivity

Tested: Human, Mouse, Rat, Pig

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

EB05284 (0.05µg/ml) staining of Human Thymus (A) and Molt4 (B) lysates (35µg total protein in RIPA buffer). Primary incubated for 1 hour. Detected by chemiluminescence.

EB05284 (0.03µg/ml) staining of Human (A) and Pig (B) Spleen lysates (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

EB05284 (0.05µg/ml) staining of Mouse (A+C) and Rat (B+D) Brain (A+B) and Spleen (C+D) lysates (35µg total protein in RIPA buffer). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.

EB05284 (1µg/ml) staining of paraffin embedded Human Lymph Node. Microwaved antigen retrieval with citrate buffer pH 6, HRP-staining.

EB05284 Negative Control showing staining of paraffin embedded Human Lymph Node, with no primary antibody.