

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB05631 - Goat Anti-GTRAP3-18 / JWA Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ARL6IP5, GTRAP3-18, JWA, ADP-ribosylation-like factor 6 interacting protein 5, DERP11, HSPC127, PRAF3, addicsin, hp22, jmx, PRA1 domain family 3, cytoskeleton related vitamin A responsive protein, dermal papilla derived protein 11, glutamate transporter EEAC1-associated protein, putative MAPK activating protein PM27

Official Symbol: ARL6IP5

Accession Number(s): NP_006398.1

Human GeneID(s): 10550

Immunogen

Peptide with sequence C-NRLTDYISKVKE, from the C Terminus of the protein sequence according to NP_006398.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 20kDa band observed in Human Brain lysates (predicted MW of 21.6kDa according to NP_006398.1). Recommended for use at 0.1-0.3µg/ml.

IHC: Paraffin embedded Human Tonsil and Small Intestine. Recommended concentration: 3.75µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

EB05631 staining (0.1μg/ml) of Human Brain lysate (RIPA buffer, 35μg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.

EB05631 (3.75μg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

EB05631 (3.75µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.