

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10176 - Goat Anti-HCN3 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: hyperpolarization activated cyclic nucleotide-gated potassium channel

3, KIAA1535, MGC131493, potassium/sodium hyperpolarization-activated cyclic

nucleotide-gated channel 3, HCN3

Official Symbol: HCN3

Accession Number(s): NP_065948.1

Human GenelD(s): 57657

Non-Human GenelD(s): 15168 (mouse)

Immunogen

 $\label{problem} \mbox{Peptide with sequence C-RGVRGRAPSTGAQ, from the internal region of the protein}$

sequence according to NP_065948.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

 $\label{lem:chromatography using the immunizing peptide.} \\$

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 85kDa band observed in Rat Kidney lysates (calculated MW of

86.0 kDa according to Human NP_065948.1 and 86.8 kDa according to Rat NP_446137.1).

Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

250kDa 150kDa 100kDa 75kDa

50kDa

37kDa

25kDa 20kDa

15kDa

EB10176 (1 μ g/ml) staining of Rat Kidney lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.