

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB05586 - Goat Anti-ICSBP1 / IRF8 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: IRF8, ICSBP1, interferon consensus sequence binding protein 1, IRF-8, H-ICSBP, interferon regulatory factor 8 Official Symbol: IRF8 Accession Number(s): NP_002154.1 Human GeneID(s): <u>3394</u> Non-Human GeneID(s): 15900 (mouse)

Immunogen

Peptide with sequence C-QRSFFRENQQITV, from the C Terminus of the protein sequence according to NP_002154.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serun albumin.


Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:8000.
Western blot: Approx 45-50kDa band seen in K562 lysate (Predicted MW of 47kDa according to NP_002154.1). Recommended concentration: 0.5-2µg/ml.
IHC: Paraffin embedded Human Tonsil. Recommended concentration: 5µg/ml.

Species Reactivity

Tested: Human Expected from sequence similarity: Human, Mouse, Rat, Cow



EB05586 staining (1µg/ml) of K562 lysate (RIPA buffer, 30µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



EB05586 (5µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.