

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09708 - Goat Anti-IKAROS / IKZF1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** IKZF1, IKAROS family zinc finger 1 (Ikaros), Hs.54452, IK1, IKAROS, LYF1, PRO0758, ZNFN1A1, hlk-1, CLL-associated antigen KW-6, Ikaros (zinc finger protein), zinc finger protein, subfamily 1A, 1 (Ikaros), lymphoid transcription factor LyF-1

**Official Symbol:** IKZF1

**Accession Number(s):** NP\_006051.1

**Human GeneID(s):** [10320](#)

### Immunogen

Peptide with sequence C-DTPYDSSASYEKEN, from the internal region of the protein sequence according to NP\_006051.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:1000.

**Western blot:** Approx 65kDa band observed in lysates of cell line MOLT4 (calculated MW of 57.5kDa according to NP\_006051.1). Recommended concentration: 1-3µg/ml. Similar results were obtained with EB07847, which was developed against a different part of the protein.

**IHC:** In paraffin embedded Human Tonsil shows textured staining in some nuclei of the lymphoid tissue. Recommended concentration, 2.5-3.8µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Dog, Cow

EB09708 (1µg/ml) staining of MOLT4 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.

EB09708 (3.8µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer  
pH 6, AP-staining.