

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09138 - Goat Anti-ITM2B Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** ITM2B, integral membrane protein 2B, ABRI, BRI, BRI2, BRICD2B, E25B, E3-16, FBD, BRICHOS domain containing 2B, OTTHUMP00000018394

**Official Symbol:** ITM2B

**Accession Number(s):** NP\_068839.1

**Human GeneID(s):** [9445](#)

**Non-Human GeneID(s):** 16432 (mouse), 290364 (rat)

### Immunogen

Peptide with sequence C-HDKETYKLQRRETIK, from the internal region of the protein sequence according to NP\_068839.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:8000.

**Western blot:** Approx. 33kDa band observed in Human Brain (Cerebellum and Hippocampus) lysates (calculated MW of 30.3kDa according to NP\_068839.1).  
Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Small Intestine. Recommended concentration: 2.5µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Cow

EB09138 (1µg/ml) staining of Human Hippocampus lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

EB09138 (2.5µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.