

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

**Enquiries:** 

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08090-B - Goat Anti-KCC3 / SLC12A6, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** SLC12A6, solute carrier family 12 (potassium/chloride transporter), member 6, ACCPN, KCC3, KCC3A, KCC3B, K-CI cotransporter 3, electroneutral potassium-chloride cotransporter 3, potassium chloride cotransporter 3, potassium chloride cotransporter KCC3a-S3, potassium-chloride transporter-3a, potassium-chloride transporter-3b, solute carrier family 12 (potassium/chloride transporters), member 6, DKFZP434D2135

Official Symbol: SLC12A6

Accession Number(s): NP\_598408.1; NP\_005126.1; NP\_001035959.1;

NP\_001035961.1; NP\_001035962.1

Human GenelD(s): 9990

Important Comments: This antibody is expected to recognise all reported isofoms (NP\_598408.1; NP\_005126.1; NP\_001035959.1; NP\_001035961.1; NP\_001035962.1); Reported variants represent identicial protein: NP\_001035959.1; NP\_001035960.1

#### **Immunogen**

Peptide with sequence SRHIDVCSKTKE., from the internal region of the protein sequence according to NP\_598408.1; NP\_005126.1; NP\_001035959.1; NP\_001035961.1; NP\_001035962.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

**Western blot:** Approx 120kDa band observed in Human Heart lysates (calculated MW ranging 121-128kDakDa according to NP\_598408.1; NP\_005126.1; NP\_001035959.1; NP\_001035960.1; NP\_001035961.1; NP\_001035962.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

Biotinylated EB08090 (0.1µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.