

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05469 - Goat Anti-KLF8 / BKLF3 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** KLF8, BKLF3, Kruppel-like factor 8, RP13-1021K9.1, DKFZp686O08126, DXS741, MGC138314, ZNF741, zinc finger protein 741, basic kruppel-like factor 3

**Official Symbol:** KLF8

**Accession Number(s):** NP\_009181.2; NP\_001152768.1

**Human GeneID(s):** [11279](#)

**Important Comments:** This antibody is expected to recognise isoform 1 (NP\_009181.2) and isoform 2 (NP\_001152768.1).

### Immunogen

Peptide with sequence VDMDKLINNLEVQ-C, from the N Terminus of the protein sequence according to NP\_009181.2; NP\_001152768.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 50kDa band observed in Human Kidney lysates (calculated MW of 39.3 according to NP\_009181.2). In glioblastomas this protein was reported to run at approx. 60kDa (Ding et al. J Biol Chem. 2005 Feb 25;280(8):6802-15; PMID: 15557280).


Recommended concentration: 1-3µg/ml.

**IHC:** Paraffin embedded Human Small Intestine. Recommended concentration: 3.75µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Pig



EB05469 staining (2 $\mu$ g/ml) of Human Kidney lysate (RIPA buffer, 30 $\mu$ g total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.



EB05469 (3.75 $\mu$ g/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.