

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06418-T - Goat Anti-LAT1 Antibody - Trial

Size: 20µg specific antibody in 40µl



### Target Protein

**Principal Names:** LAT1, LAT, linker for activation of T cells, pp36, 36 kDa  
phospho-tyrosine adaptor protein, linker for activation of T-cells family member 1

**Official Symbol:** LAT

**Accession Number(s):** NP\_055202.1; NP\_001014987.1; NP\_001014989.1;  
NP\_001014988.1

**Human GeneID(s):** [27040](#)

**Important Comments:** This antibody is expected to recognize isoform a (NP\_055202.1),  
isoform b (NP\_001014987.1 and NP\_001014989.1) and isoform c (NP\_001014988.1).

### Immunogen

Peptide with sequence C-GAPDYENLQELN, from the C Terminus of the protein  
sequence according to NP\_055202.1; NP\_001014987.1; NP\_001014989.1;  
NP\_001014988.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity  
chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum  
albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:2000.

**Western blot:** Approx 37kDa band observed in Human Peripheral Blood  
Mononucleocytes lysates (calculated MW of 27.9kDa according to NP\_055202.1). The  
observed molecular weight corresponds to earlier findings with different antibodies from  
other commercial sources. Recommended concentration: 0.01-0.03µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat



EB06418 (0.01 $\mu$ g/ml) staining of Peripheral Blood Mononucleocytes lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.