

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07538 - Goat Anti-MBD2 (isoform 1) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** MBD2, methyl-CpG binding domain protein 2, DKFZp586O0821, DMTase, NY-CO-41

**Official Symbol:** MBD2

**Accession Number(s):** NP\_003918.1

**Human GeneID(s):** [8932](#)

**Non-Human GeneID(s):** 17191 (mouse)

**Important Comments:** This antibody is expected to recognize isoform 1 (NP\_003918.1) only.

### Immunogen

Peptide with sequence C-RNDPLNQNKGPDLN, from the internal region of the protein sequence according to NP\_003918.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:64000.

**Western blot:** Approx 48-50kDa band observed in lysates of cell line Jurkat (calculated MW of 43.3kDa according to NP\_003918.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Tan et al, Mol Cell Biol. 2006 Oct;26(19):7224-35.; PMID: 16980624). Recommended concentration: 0.3-1µg/ml.

**IHC:** In paraffin embedded Human Colon shows nuclear staining of enterocytes. Recommended concentration, 2-4µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog

### Specific Reference

**This antibody has been successfully used (in WB on HeLa (46kDa and 27kDa)) in the following paper:**

Spruijt CG, Bartels SJ, Brinkman AB, Tjeertes JV, Poser I, Stunnenberg HG, Vermeulen M.


CDK2AP1/DOC-1 is a bona fide subunit of the Mi-2/NuRD complex.

Mol Biosyst. 2010 Sep;6(9):1700-6.

PMID: 20523938



EB07538 (0.3 $\mu$ g/ml) staining of Jurkat cell lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.



EB07538 (2.5 $\mu$ g/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.