

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10286 - Goat Anti-MGAT1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: GLCNAC-TI, GLCT1, GLYT1, GNT-1, GNT-1, mannosyl (alpha-1,3-)-glycoprotein beta-1,2-N-acetylglucosaminyltransferase, MGAT, N-glycosyl-oligosaccharide-glycoprotein N-acetylglucosaminyltransferase, OTTHUMP00000161546, MGAT1 Official Symbol: MGAT1 Accession Number(s): NP_002397.2 Human GeneID(s): 4245 Non-Human GeneID(s): 17308 (mouse), 81519 (rat) Important Comments: Reported variants represent identical protein: NP_002397.2, NP_001108090.1, NP_001108091.1, NP_001108092.1, NP_001108089.1.

Immunogen

Peptide with sequence C-QVEKVRTNDR, from the internal region of the protein sequence according to NP_002397.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: In transfected HEK293 transiently expressing full-length Human MGAT1 (myc and DYKDDDDK tagged), a band of approx. 55kDa was observed. No bands were observed in mock-transfected HEK293 and the same band was observed using anti-myc tag antibody. Recommended concentration, $0.2-1\mu g/ml$.

Purchase the +ve control! The transfected lysate used to QC this product is available for sale. Please email sales@everestbiotech.com to request datasheet, pricing and delivery information.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

HEK293 lysate (10ug protein in RIPA buffer) overexpressing Human MGAT1 with C-terminal MYC tag probed with EB10286 (1ug/ml) in Lane A and probed with anti-MYC Tag (1/1000) in lane C. Mock-transfected HEK293 probed with EB10286 (1mg/ml) in Lane B. Primary incubations were for 1 hour. Detected by chemiluminescence.