

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05055 - Goat Anti-MTR Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: MTR, 5-methyltetrahydrofolate-homocysteine methyltransferase, 5-methyltetrahydrofolate-homocysteine methyltransferase 1, FLJ33168, FLJ43216, FLJ45386, MS, cblG, cobalamin-dependent methionine synthase, methionine synthase, vitamin-B12 dependent methionine synthase

Official Symbol: MTR

Accession Number(s): NP_000245.2

Human GeneID(s): [4548](#)

Non-Human GeneID(s): 81522 (rat)

Immunogen

Peptide with sequence C-VEKWLGPILGYDTD, from the C Terminus of the protein sequence according to NP_000245.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

IHC: In paraffin embedded Human Adrenal Gland shows cytoplasm staining in the cortex. Recommended concentration: 3-5µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig

EB05055 (3.8µg/ml) staining of paraffin embedded Human Adrenal Gland. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.