

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07870 - Goat Anti-c-Myb (aa281-294) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: c-myb, MYB, v-myb myeloblastosis viral oncogene homolog (avian), c-myb_CDS, efg, Avian myeloblastosis viral (v-myb) oncogene homolog, c-myb protein (140 AA), c-myb10A_CDS, c-myb13A_CDS, c-myb14A_CDS, c-myb8B_CDS, v-myb avian myeloblastosis viral oncogene homolog, v-myb myeloblastosis viral oncogene homolog

Official Symbol: MYB

Accession Number(s): NP_001123645.1; NP_005366.2; NP_001123644.1; NP_001155128.1; NP_001155129.1; NP_001155130.1; NP_001155131.1

Human GeneID(s): [4602](#)

Non-Human GeneID(s): 17863 (mouse)

Important Comments: This antibody is expected to recognize isoforms 1-7 only (Numbering from NCBI RefSeq). (NP_001123645.1; NP_005366.2; NP_001123644.1; NP_001155128.1; NP_001155129.1; NP_001155130.1; NP_001155131.1)

Immunogen

Peptide with sequence C-QRHYNDEDPEKEKR, from the internal region of the protein sequence according to NP_001123645.1; NP_005366.2; NP_001123644.1; NP_001155128.1; NP_001155129.1; NP_001155130.1; NP_001155131.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:1000.

Western blot: Approx 70kDa band observed in Human Colon lysates (calculated MW of 72.3kDa according to NP_005366.2). Recommended concentration: 0.5-2µg/ml.

IHC: In paraffin embedded Kidney shows nuclear staining in all cells. Recommended concentration: 2-4µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog

EB07870 (xµg/ml) staining of Human Colon lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.

EB07870 (2µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining.