

### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09020 - Goat Anti-MYH9 Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** myosin, heavy chain 9, non-muscle, RP1-68O2.1, DFNA17, EPSTS, FTNS, MGC104539, MHA, NMHC-II-A, NMMHCA, MYH9 variant protein, OTTHUMP00000028706, cellular myosin heavy chain, type A, myosin, heavy polypeptide 9, non-muscle, non-muscle myosin heavy chain, non-muscle myosin heavy polypeptide 9,

Official Symbol: MYH9

Accession Number(s): NP\_002464.1

nonmuscle myosin heavy chain II-A

Human GeneID(s): 4627

## Immunogen

Peptide with sequence C-DQINTDLNLERSH, from the internal region of the protein sequence according to NP\_002464.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 250kDa band observed in lysates of Human Peripheral Blood Mononucleocytes and of NIH3T3 (calculated MW of 227kDa according to Human NP\_002464.1 and to Mouse NP\_071855.2). Recommended concentration: 0.3-1μg/ml.

### **Species Reactivity**

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog

EB09020 (0.3µg/ml) staining of Human peripheral blood lymphocytes lysate (35µg protein in RIPA buffer).

Primary incubation was 1 hour. Detected by chemiluminescence.

EB09020 (1 $\mu$ g/ml) staining of NIH3T3 lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.