

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11023 - Goat Anti-NMNAT3 (aa149-162) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NMNAT3, nicotinamide nucleotide adenylyltransferase 3, PNAT3, NMN adenylyltransferase 3, NaMN adenylyltransferase 3, nicotinamide mononucleotide adenylyltransferase 3, nicotinate-nucleotide adenylyltransferase 3, pyridine nucleotide adenylyltransferase 3, NaMN adenylyltransferase 3, nicotinamide mononucleotide adenylyltransferase 3, nicotinamide nucleotide adenylyltransferase 3, nicotinate-nucleotide adenylyltransferase 3, NMN adenylyltransferase 3, PNAT3, pyridine nucleotide adenylyltransferase 3, NMNAT3

Official Symbol: NMNAT3

Accession Number(s): NP_835471.1; NP_001186976.1

Human GenelD(s): [349565](#)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_835471.1; NP_001186976.1).

Immunogen

Peptide with sequence C-QHNIHLAKEPVQNE, from the internal region of the protein sequence according to NP_835471.1; NP_001186976.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 26kDa band observed in Human Kidney lysates (calculated MW of 24.1kDa according to NP_835471.1). Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog



EB11023 (0.1 μ g/ml) staining of Human Kidney lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.