

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07246 - Goat Anti-NONO / p54NRB Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NONO, P54NRB, non-POU domain containing, octamer-binding, HGNC:7871, NMT55, NRB54, P54, Nuclear RNA-binding protein, 54-kD, non-POU-domain-containing, octamer-binding, non-Pou domain-containing octamer (ATGCAAAT) binding protein

Official Symbol: NONO

Accession Number(s): NP_031389.3

Human GeneID(s): [4841](#)

Non-Human GeneID(s): 53610 (mouse), 317259 (rat)

Immunogen

Peptide with sequence C-NRAAPGAEFAPNK, from the C Terminus of the protein sequence according to NP_031389.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 60kDa band observed in Mouse Brain lysates (calculated MW of 54.2kDa according to Human NP_031389.3 and 54.5kDa according to Mouse NP_075633.2). In transfected HEK293 transiently expressing Human NONO a band of approx. 55kDa is observed. This band is not observed in the non-transfected HEK293. The endogenous 50kDa protein is observed in both. Recommended concentration: 0.1-0.3µg/ml.

IHC: In paraffin embedded Human Spleen shows nuclear staining in select splenocytes. Recommended concentration: 3-6µg/ml. Paraffin embedded Human Testis. Recommended concentration: 3.75µg/ml.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog

EB07246 (0.1µg/ml) staining of Mouse Brain lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.

HEK293 overexpressing Human NONO (RC206688) and probed with EB07246 (mock transfection in first lane),
tested by Origene.

EB07246 (3.8µg/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer
pH 6, AP-staining.

EB07246 (3.75µg/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer
pH 6, AP-staining.