

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06087 - Goat Anti-NUMBL Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** NUMBL, numb homolog (Drosophila)-like, NBL, CAG3A, CTG3a, NUMBR, NUMB-R, TNRC23, numb (Drosophila) homolog-like, NUMBLIKE, TNRC23, numb homolog-like

**Official Symbol:** NUMBL

**Accession Number(s):** NP\_004747.1

**Human GeneID(s):** [9253](#)

**Non-Human GeneID(s):** 18223 (mouse)

### Immunogen

Peptide with sequence SRSAAASGGPRRPE-C, from the N Terminus of the protein sequence according to NP\_004747.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** In transfected HEK293 transiently expressing Human Numbl and DYKDDDDK tagged), a band of approx. 85kDa is observed. No bands are observed in mock-transfected HEK293 and the same band is observed using anti-myc antibody. Recommended concentration, 0.5-1µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat



HEK293 lysate (10ug protein in RIPA buffer) overexpressing Human Numbl with C-terminal MYC tag probed with EB06087 (1ug/ml) in Lane A and probed with anti-MYC Tag (1/1000) in lane C. Mock-transfected HEK293 probed with EB06087 (1mg/ml) in Lane B. Primary incubations were for 1 hour. Detected by chemiluminescence.