

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08608 - Goat Anti-OX1R and OX2R Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: OX1R, OX2R, HCRTR1, hypocretin (orexin) receptor 1, hypocretin receptor 1, hypocretin receptor 1, orexin receptor 1, orexin receptor-1, HCRTR2, hypocretin (orexin) receptor 2, hypocretin receptor 2, hypocretin receptor-2, orexin

receptor 2, orexin receptor-2

Official Symbol: HCRTR1

Accession Number(s): NP_001516.2; NP_001517.2

Human GenelD(s): 3061, 3062

Non-Human GeneID(s): 230777 (mouse), 25593 (rat)

Important Comments: This antibody is expected to recognise both the human proteins

OX1R and OX2R.

Immunogen

Peptide with sequence C-YNFLSGKFREQFK, from the internal region of the protein sequence according to NP_001516.2; NP_001517.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 48kDa band observed in Human Brain (Frontal Cortex), Mouse Brain and Rat Brain lysates (calculated MW of 47.5kDa according to NP_001516.1 and NP_001517.1). Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

EB08608 (0.5μg/ml) staining of Human Frontal Cortex lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.

EB08608 (1µg/ml) staining of Mouse Brain lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.