

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05880-B - Goat Anti-PARK7 / DJ-1, biotinylated Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PARK7, parkinson protein 7, DJ-1, DJ1, HEL-S-67p, Parkinson disease (autosomal recessive, early onset) 7, epididymis secretory sperm binding protein Li 67p, oncogene DJ1, protein DJ-1, CTA-215D11.1, FLJ27376, FLJ34360, FLJ92274

Official Symbol: PARK7

Accession Number(s): NP_009193.2

Human GeneID(s): [11315](#)

Non-Human GeneID(s): 57320 (mouse), 117287 (rat)

Important Comments: Reported variants represent identical protein: NP_009193.2, NP_001116849.1

Immunogen

Peptide with sequence C-AAQVKAPLVLKD., from the C Terminus of the protein sequence according to NP_009193.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 24kDa band observed in lysates of cell line Jurkat (calculated MW of 19.9kDa according to NP_009193.2). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat



Biotinylated EB05880 (0.2µg/ml) staining of Jurkat lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.