

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07477-B - Goat Anti-PEBP1 / RKIP (Internal), Biotinylated Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PEBP1, phosphatidylethanolamine binding protein 1, HCNP, HCNPPp, HEL-210, HEL-S-34, HEL-S-96, PBP, PEBP, PEBP-1, RKIP, Raf kinase inhibitory protein, epididymis luminal protein 210, epididymis secretory protein Li 34, epididymis secretory protein Li 96, hippocampal cholinergic neurostimulating peptide, neuropolypeptide h3, prostatic binding protein, raf kinase inhibitor protein

Official Symbol: PEBP1

Accession Number(s): NP_002558.1

Human GeneID(s): [5037](#)

Non-Human GeneID(s): 23980 (mouse), 29542 (rat)

Immunogen

Peptide with sequence C-DPDAPSRKDPKYRE., from the internal region of the protein sequence according to NP_002558.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 23kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 21.1kDa according to NP_002558.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



Biotinylated EB07477 (0.1µg/ml) staining of Human Cerebellum lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.