

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB11523 - Goat Anti-PLK1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: cell cycle regulated protein kinase, OTTHUMP00000122462, PLK, PLK1, PLK-1, polo (Drosophia)-like kinase, polo like kinase, polo-like kinase 1, serine/threonine-protein kinase 13, serine/threonine-protein kinase PLK1, STPK13

Official Symbol: PLK1

Accession Number(s): NP_005021.2

Human GeneID(s): 5347

Immunogen

Peptide with sequence TSGYIPARLPITC, from the internal region of the protein sequence according to NP_005021.2.

Please note the <u>peptide</u> is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 65kDa band observed in lysates of cell line K562 (calculated MW of 68.3kDa according to NP_005021.2). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa

15kDa

EB11523 ($1\mu g/ml$) staining of K562 lysate ($35\mu g$ protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.